



Nerve repair, made SMRT™

Remplir is engineered with Orthocell's proprietary SMRT™ technology to help patients reclaim touch, movement, and their lives.^{1,2}



Remplir™

Peripheral nerve repair:

A smarter way to support nerve healing

The best predictor of success in peripheral nerve surgery is a tension-free, well-aligned coaptation using minimal epineurial microsutures to optimize axonal regeneration.³

Yet depending on the type and location of the repair, achieving this can be challenging and outcome predictability has changed little over the past decades.³⁻⁵

Biomaterial choice may play a larger role than previously thought.


Many materials fail to fully integrate with native tissue, leaving residues that impede healing. Rigid materials can also be difficult to handle and complicate management of size mismatches.^{6,7}

Remplir is engineered with Orthocell's proprietary SMRT™ technology to:

- mimic the natural epineurium with an absolute collagen wrap¹
- create an optimal healing environment for nerve regeneration^{1,2}
- be easy to handle and suture-sparing, simplifying the surgical process^{1,2}
- protect the repair site from scarring, adhesions, and inflammation.¹


Repair with Remplir:

One wrap, multiple applications



Connect

Enables tensionless repair of transected nerve injuries, supporting direct repairs with or without grafts, as well as nerve transfers and free functioning muscle transfers.^{1,2}



Protect


Helps manage non-transected nerve injuries by reducing the risk of adhesion formation and supporting smooth nerve gliding in conditions such as carpal and cubital tunnel syndrome.¹



Cap

Prevents neuroma regrowth after resection of a terminal neuroma, or formation of new neuromas post amputation.¹

Protects, integrates and provides a template for restoration of native epineurium



Benefits of naturally occurring collagen structure

1 Smooth outer surface

- Densely packed collagen bundles
- Cell occlusive surface minimizes the risk of adhesions and facilitates nerve gliding¹
- Permits passage of fluids and nutrients while retaining neurotrophic factors.¹


2 Rough inner surface

- Loosely distributed collagen bundles
- Creates favorable microenvironment for nerve healing.^{1,2}

Remplir pre-clinical evidence⁸




Remplir-assisted repair: restored nerve structure




Intact rat sciatic nerve

- Longitudinally-aligned nerve fibers surrounded by protective outer layer of epineurium



Remplir-wrapped rat sciatic nerve

- Restoration of nerve fiber structure
- Epineurium-like tissue is restored as Remplir is resorbed
- No scar tissue, inflammation or foreign body reaction observed
- No compression of nerve




Comparator-wrapped rat sciatic nerve

- Abnormal nerve structure encapsulated in thick, inflamed fibrous tissue (*blue*)
- Inflammation and foreign body reaction observed
- Compression of nerve fibers with nerve entrapment (*red*) and separation of epineurium from nerve.

Remplir pre-clinical evidence⁸




Facilitates gliding
and minimizes adhesion




Intact rat sciatic nerve

- Pearly white appearance
- Moves freely within surrounding soft tissue



Remplir-wrapped rat sciatic nerve

- Integrated into nerve tissue with appearance resembling pre-injured state (*green*)
- No evidence of inflammation
- Minimal adhesions or scar tissue formation




Comparator-wrapped rat sciatic nerve

- Not integrated with nerve tissue (*blue*)
- Multiple points of adhesion to surrounding tissue
- Inflammatory reaction observed


Benefits of augmenting suture repair with Remplir

Suture repair with Remplir



Remplir forms a protective, epineurial-like barrier around the repaired nerve, with no inflammatory response observed. Vascularization occurs both external to Remplir and within the nerve. Formation of adipose and connective tissue external to Remplir cushions the nerve and allows nerve gliding.


Suture repair without Remplir



No regeneration of protective epineurium or induction of adipose or connective tissue was observed after suture repair without Remplir, increasing the risk of adhesions and nerve compression.


Integration and restoration of nerve tissue

Early integration of Remplir, crucial for maintaining the structural integrity of the repaired nerve during early healing, was observed after 4 weeks.




4 weeks

Early integration of Remplir and formation of epineurium-like tissue (ELT)



12 weeks

Remplir largely resorbed, and ELT is becoming restored



24 weeks

Remplir completely resorbed and ELT is restored

As Remplir degrades over 24 weeks, the nerve is increasingly protected by epineurium-like tissue. Physiological inflammation (indicated by dark purple cellular staining) is the normal healing process observed across the observation period. No foreign body response was observed.

A complete range of nerve wraps



15x20mm



20x30mm



30x40mm



40x50mm

Nerve diameter

Up to 3mm

Up to 5mm

Up to 10mm

Used for longitudinal
wrapping

Note: Smooth side away from nerve

Order codes

ON-152

ON-203

ON-304

ON-405



About us

Orthocell Ltd, headquartered in Perth, Australia, is the home of Remplir

Since 2010, we've been developing and exporting innovative regenerative medicine products worldwide. To date, more than 70,000 patients have benefited from an Orthocell collagen medical device, with no reported adverse events as of March 2025. Every product reflects our mission: improving mobility, function, and quality of life. We collaborate with leading surgeons and centers of excellence to translate research into practical solutions that raise the standard of care. Our deep understanding of tissue engineering, particularly the interaction between cells and scaffolds, allows us to enhance the body's natural healing and regeneration processes. Orthocell products are designed to simplify procedures, save operative time, and deliver consistent, predictable outcomes, supporting surgeons in providing the best care for their patients.

References. **1.** Data on file. **2.** O'Beirne, A., J. Cullen, E. Landao-Bassonga, M. Zheng, C. Lee, P. Kaluskar, A. Tai and M. Zheng (2024). "Reconstruction of Upper Extremity Peripheral Nerve Injuries Using an Epineurial-Like Collagen Device—A Prospective Clinical Study." *Journal of Reconstructive Microsurgery Open* 09(01): e78-e88. **3.** Siemionow, M. and G. Brzezicki (2009). "Chapter 8: Current techniques and concepts in peripheral nerve repair." *Int Rev Neurobiol* 87: 141-172. **4.** Ruijs, A.C., J.B. Jaquet, S. Kalmijn, H. Giele, S.E. Hovius (2005). "Median and ulnar nerve injuries: a meta-analysis of predictors of motor and sensory recovery after modern microsurgical nerve repair." *Plast Reconstr Surg* 116(2):484-94. **5.** Braga Silva, J., G. M. Marchese, C. G. Cauduro and M. Debiasi (2017). "Nerve conduits for treating peripheral nerve injuries: A systematic literature review." *Hand Surg Rehabil* 36(2): 71-85. **6.** Aman M, Preisner F, Mayrhofer-Schmid M, et al. Inferior Long-Term Results of a Randomized Controlled Trial Initially Demonstrating Enhanced Sensory Nerve Recovery Using a Chitosan Nerve Tube. *Plast Reconstr Surg.* 2025;155(2):364-373. Doi:10.1097/PRS.00000000000011526 **7.** Zeng W, Osterman M, Stern PJ. Inflammatory Reactions to Xenogenic Nerve Wraps: A Report of Three Cases. *JBJS Case Connect.* 2019;9(3):e0302. Doi:10.2106/JBJS.CC.18.00302 **8.** Data on file, 4 weeks post-transection.

© 2025 Orthocell Ltd. All rights reserved. MK--0002094

Head Office (Australia)

P: +61 8 9360 2888

E: general@orthocell.com

W: orthocell.com

US Product Inquiries

P: 919-842-5915

E: orthocell@unipharlogistics.com

W: remplir.orthocell.com